

Available online at www.sciencedirect.com



Discrete Applied Mathematics 149 (2005) 219

## DISCRETE APPLIED MATHEMATICS

www.elsevier.com/locate/dam

## Author index to volume 149

Bloch, J.C., The complexity of modular decomposition of Boolean functions	(1-3)	1-13
Čepek, O. and P. Kučera, Known and new classes of generalized Horn formulae		
with polynomial recognition and SAT testing	(1-3)	14 - 52
Cohen, D., M. Cooper, P. Jeavons and A. Krokhin, Supermodular functions and		
the complexity of MAX CSP	(1-3)	53-72
Cooper, M., see D. Cohen	(1-3)	53-72
Cusick, T.W. and Y. Li, k-th order symmetric SAC boolean functions and bisecting		
binomial coefficients	(1-3)	73-86
Faye, A. and Q. Trinh, A polyhedral approach for a constrained quadratic 0-1		
problem	(1-3)	87 - 100
Geist, D., see B. Schieber	(1-3)	154-173
Golumbic, M.C., see A. Mintz	(1-3)	131-153
Hatami, H. and H. Maserrat, On the computational complexity of defining sets	(1-3)	101-110
Jansen, T. and I. Wegener, Real royal road functions—where crossover provably is		
essential	(1-3)	111-125
Jeavons, P., see D. Cohen	(1-3)	53-72
Krokhin, A., see D. Cohen	(1-3)	53-72
Kučera, P., On the size of maximum renamable Horn sub-CNF	(1-3)	126-130
Kučera, P., see O. Čepek	(1-3)	14-52
Li, Y., see T.W. Cusick	(1-3)	73-86
Maserrat, H., see H. Hatami	(1-3)	101-110
Mintz, A. and M.C. Golumbic, Factoring Boolean functions using graph		
partitioning	(1-3)	131-153
Schieber, B., D. Geist and A. Zaks, Computing the minimum DNF representation		
of Boolean functions defined by intervals	(1-3)	154-173
Trinh, Q. see A. Faye	(1-3)	87-100
Wegener, I., see T. Jansen	(1-3)	111-125
Wild, M., The many benefits of putting stack filters into disjunctive or conjunctive		
normal form	(1-3)	174-191
Xiao, W., Linear symmetries of Boolean functions	(1-3)	192-199
Zaks, A., see B. Schieber	(1-3)	154-173
Zverovich, I.E., Characterizations of closed classes of Boolean functions in terms of		
forbidden subfunctions and Post classes	(1-3)	200-218